

Problems: unreported catch

- a. night
- b. launch
- c. cabins

F0.1

- a. m
- b. selectivity

Hooking mortality

Sex

Safe harvest level.

- a. remove hooking mortality
- b. tribal harvest

creel survey

1. check previous years format to ensure it is the same.
2. What is going on with the low trawl catches.
3. resolve what is going on with the

/autofs/users/bioram/myers/walleye/minn/jan2000/MLGNet99.v6x

Here's more data. The first file, MLAgeHrT.v6x, gives tribal harvest by age for both the spring and fall seasons. This can replace the file I sent to you earlier (MLAgeHrv) that just had spring tribal harvest by age. Note that the fall tribal harvest by age estimates used aging results from MN DNRs gill net surveys. The second file, MLFshFal.v6x, has a record for each walleye harvested during the fall by the tribes.

Ram,

Here's some additional Mille Lacs assessment data from Rick Bruesewitz. This includes data from 8 newer assessment gill nets, in addition to the standard 32-net assessment, and the trawl results are also included.

"MLGnTr99.v6x")

Ram,

I received 1999 Mille Lacs gill net assessment data from MN DNR's original 32 nets today from Rick B. Here 'tis.

Rick

[Part 2, Application/OCTET-STREAM (Name: "MLGNet99.v6x") 39KB]

Rick

if survey="Original gill net assessment";